NOCOSPRAY USER MANUAL



CAREFULLY READ THE ENTIRE USER'S MANUAL THAT COMES WITH THE NOCOSPRAY ROBOT!



Use gloves and eye protection when handling any chemicals used for the robot.



Check that the bottle contains sufficient amount of chemistry. Mark the level *before* the procedure and check after.




Secure the spray nozzle onto the bottle.

Keep the cap!



Attach the spray nozzle with the bottle in place on the Nocospray robot. Ensure the nozzle is firmly attached.



Place the Nocospray robot on a stable surface. Choose a corner for optimal effect.



Nocolyse OneShot: room volume (m³) x3 Nocolyse Neutral: room volume (m³) x1

Set the correct room volume. The setting depends on the size of the room and the type of chemistry used.



Press start in the center of the dial. The robot will beep for 15 seconds and start automatically.

eave the room!



Nocolyse OneShot: 2 hours contact time Nocolyse Neutral: 30 min. contact time

The room must remain closed/locked during diffusion and throughout the contact time. Hang a poster on the door with a warning about "No entry!"



Detach the bottle and replace the cap. Rinse the spray nozzle and tube under water. Click the nozzle back into place in the robot.



Adresse: Telefon: E-mail: Web:

Teglverksveien 47, 3057 Solbergelva 32 83 59 00 info@bragemedical.no www.bragemedical.no